PCT

WORLD INTELLECTUAL PROP International I

INTERNATIONAL APPLICATION PUBLISHED UNDE

(51) International Patent Classification 6:

C12Q 1/66

(11) I

96077591

(43) International Publication Date:

14 March 1996 (14.03.96)

(21) International Application Number:

PCT/GB95/02085

A1

(22) International Filing Date:

4 September 1995 (04.09.95)

(30) Priority Data:

9417793.8

5 September 1994 (05.09.94)

(71) Applicant (for all designated States except US): CELSIS INTERNATIONAL PLC [GB/GB]; Science Park, Milton Road, Cambridge CB4 4FX (GB).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): FOOTE, Nicholas, Peter, Martin [GB/GB]: 15 The Woodlands, Linton, Cambridge CB1 6UF (GB). NOBLE, Michael [GB/GB]; 41 River Lane, Cambridge CB5 8HP (GB). GRANT, Peter, Leonard [GB/GB]; 28 Russett Avenue, Needingworth, Cambs PE17 3UE (GB). EVANS, Christopher, Thomas [GB/GB]; Stable Cottage, Fowlmere Road, Heydon, Hertfordshire SG8 8PU (GB).
- (74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, ČN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT. SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: BIOLUMINESCENCE AND ASSAY REAGENTS

(57) Abstract

A kit of two more containers, suitable for use in the estimation of ATP released from microorganisms, contains luciferin, luciferase, a reconstitution medium and, if desired, a surfactant or other cell lytic agent, and additionally contains a medium-chain alkanoic acid.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

		GB	United Kingdom	MR	Mauritania
AT	Austria	GE	Georgia	MW	Malawi
AU	Australia		Guines	NE	Niger
BB	Barbados	GN		NL	Netherlands
BE	Belgium	GR	Greece	NO	Norway
BF	Burkina Faso	HU	Hungary		•
BG	Bulgaria	1E	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy .	PL	Poland
BR	Brazil .	JP	Japan	PT	Portugal
	Belarus	KE	Kenya	RO	Romania
BY		KG	Kyrgysian	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic	, Ad	of Korea	SE	Sweden
CG	Congo	***	••	SI	Slovenia
CH	Switzerland	KR	Republic of Korea	SK	Slovakia
CI	Côte d'Ivoire	KZ	Kazakhstan	SN	Senegal
CM	Cameroon	IJ	Liechtenstein		~
CN	China	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
CZ	Czech Republic	LV	Larvia	TJ	Tajikistan
DE	Germany	MC	Monaco	TT	Trinidad and Tobago
		MD	Republic of Moldova	UA	Ukraine
DK	Denmark	MG	Madagascar	US	United States of America
ES	Spain	-	Mali	UZ.	Uzbekistan
FI	Finland	ML		VN	Viet Nam
FR	France	MN	Mongolia	4.14	

WO 96/07759 PCT/GB95/02085

BIOLUMINESCENCE AND ASSAY REAGENTS

This invention relates to a method for detecting microorganisms.

Live microorganisms can be detected with great sensitivity by measuring their ATP using a reagent based on the firefly luciferase reaction. This ATP must be extracted by lysing the microorganisms. There are presently four main ways of doing this, i.e. extracting with boiling buffer or with detergent, or using solvents or acids.

The detergent method is understandably the most popular because it involves a minimum of sample handling, is almost instantaneous, and allows a simple two-step assay, comprising: (1) adding an extractant to the sample; and (2) adding reagent and measuring light. This method is the subject of GB-A-1604249.

The detergent often actually stimulates the initial light production, but also rapidly inactivates the enzyme. Thus the light output falls to nothing over a period of a few minutes. Therefore, despite the initial stimulation, there may be a drastic loss of assay sensitivity, especially if the light output is not measured straight away.

It is known that non-ionic detergents protect, at least in part, from this inactivation. However, they may also protect bacteria from the lytic effects of the cationic detergent; therefore, a "mixed" detergent extractant is not efficient. Furthermore, non-ionic detergents themselves have a slow inactivating effect on luciferase and cannot be included as part of a stable reagent.

The only way to use a non-ionic detergent as a protectant is to add it between the additions of extractant and reagent. The assay now requires three steps, and is therefore less convenient and more expensive to automate.

An object behind the present invention is to provide a reagent having a low decay rate, i.e. <5% per minute,

10

15

20

25

30

35

WO 96/07759 PCT/GB95/02085

2

e.g. even after the addition of a detergent-based extractant, so that it can be used in a two-step assay. This could be used in even the simplest luminometer, e.g. a camera-type microtitre plate system, and would bring all the advantages of improved accuracy and sensitivity of a true low-decay rate assay.

Surprisingly, it has been found that medium chain-length fatty acids, e.g. C₅₋₁₅ alkanoic acids, are efficient luciferase protectants. Their esters or other derivatives/analogues may also be used. For example, a reagent containing octanoic acid retains low-decay rate kinetics even in the presence of an efficient extractant based on cationic detergents. It has little effect on the storage stability of luciferase and can even be added to a luciferase-luciferin reagent before it is freeze-dried, although the, say, octanoic acid is preferably included in the reagent reconstitution buffer. Such a reagent could still be used after 8 hours at 25°C or 1 week at 4°C.

The following Examples illustrate the invention.

20 Example 1

5

10

15

25

30

35

Vials of a commercially-available freeze-dried reagent preparation containing firefly luciferase, D-luciferin and stabilisers, a low decay rate cocktail, were dissolved in 5 ml of buffer without and with added octanoic acid. The latter comprised 50 mM Tris-Hepes, 9 mM magnesium acetate, 1 mM EDTA, 1 g/l octanoic acid and 0.02% sodium azide. After incubation at 15°C for 1 hour, the two preparations were compared in an assay of ATP in the presence of a bacterial lysing reagent containing cationic detergents.

The assay was performed as follows using a Berthold Biolumat LB9500T luminometer: 0.1 ml of the lysing agent was placed in a cuvette, 0.1 ml of 2 x 10⁻⁹ M Na₂ATP in sterile water was added, and the cuvette was equilibrated at 25°C for 5 minutes. 0.1 ml of the reagent was then added. The initial light output was measured by a 10 second integration, and the light signal was continuously monitored on a chart recorded so that the decay rate could

be calculated. The stability of the reconstituted reagents was compared by storage at 25°C and 4°C. The results were as follows:

	Without octanoic acid	With octanoic acid
Response to ATP (RLU/attomol)	0.62	0.42
Decay rate (% min ⁻¹)	27	<3
Stability 24 hr 25°C (%)	87	60
Stability 1 week 4°C (%)	84	76

These results show that the presence of octanoic acid provides adequate stability in solution for most applications.

15 Example 2

5

10

20

25

Vials of a high sensitivity cocktail were reconstituted and assayed as above, except that the concentration of the Na₂ATP solution was 2 x 10⁻¹⁰ M. The decay of the light signal was slower in the presence of octanoic acid, and displayed single exponential kinetics. By determining the half-life of decay and multiplying by the initial light output, an accurate calculation of the total light emitted by the reaction mixture (in arbitrary units) could be made. This was shown to be a more accurate measure of the quantity of ATP than the initial light output alone.

PCT/GB95/02085 WO 96/07759

<u>CLAIMS</u>

- A kit of two more containers, suitable for use in the estimation of ATP released from microorganisms, containing luciferin, luciferase, a reconstitution medium and, if 5 desired, a surfactant or other cell lytic agent, which additionally contains a medium-chain alkanoic acid.
 - A kit according to claim 1, wherein one container contains the medium and the acid.
- A kit according to claim 1 or claim 2, wherein the acid is octanoic acid. 10
 - A kit according to any preceding claim, wherein one container contains a freeze-dried preparation including luciferin and luciferase.
- A kit according to any preceding claim, wherein the medium is a buffer having substantially the optimum pH for 15 the luciferin-luciferase reaction.
 - Any one of the containers defined in any of claims 1 to 5 that contains the acid.
- An assay of cells, which comprises lysing the cells and measuring the ATP released by means of bioluminescence 20 in the presence of a medium-chain alkanoic acid.

INTERNATIONAL SEARCH REPORT

In onal Application No PCT/GB 95/02085

A CLASSII	SIGNATION OF SURJECT MATTER			
IPC 6	FICATION OF SUBJECT MATTER C12Q1/66			
A	o International Patent Classification (IPC) or to both national classi	fication and IPC		
	SEARCHED			
	ocumentation searched (classification system followed by classificat	on symbols)		
IPC 6	C12Q			
Documentati	con searched other than minimum documentation to the extent that	such documents are included in the fields se	arched	
Electronic d	ata base consulted during the international search (name of data base	se and, where practical, search terms used)		
	MENTS CONSIDERED TO BE RELEVANT		Relevant to claim No.	
Category *	Citation of document, with indication, where appropriate, of the r	elevant passages	Rejevant to claim No.	
	TOURNAL OF BIOLOGICAL CUENTCIDY		1-7	
٨	JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 268, no. 33, 25 November 19	SI UM EB	1-7	
	pages 24734-24741,	33 110 03,		
	W. A. FRANCISCO ET AL. 'Interac	tion of	•	
	bacterial luciferase with aldehy	de		
	substrates and inhibitors.			
	see the whole document			
		-/		
1				
ļ				
X Fw	rther documents are listed in the continuation of box C.	Y Patent family members are listed	in annex.	
* Special c	alegones of cited documents:	"I" later document published after the mi	ernational filing date	
	ment defining the general state of the art which is not	or priority date and not in conflict w cited to understand the principle or t	ith the application out	
1	idered to be of particular relevance r document but published on or after the international	invention "X" document of particular relevance; the	claimed invention	
filing	date ment which may throw doubts on priority claim(s) or	cannot be considered novel or canno involve an inventive step when the d	t be considered to	
white	ment which may drow doubt on phonic dain(s) or b is gred to establish the publication date of another ion or other special reason (as specified)	"Y" document of particular relevance; the	daimed invention	
"O" docum	ment referring to an oral disclosure, use, exhibition or	cannot be considered to involve an a document is combined with one or in ments, such combination being obvious	nore other such docu-	
	r means ment published prior to the international filing date but	in the art.		
	than the priority date claimed	"A" document member of the same pater		
Date of th	e actual completion of the international search	Date of mailing of the international s	earch report	
1 .	9 January 1996	1 6, 01, 96		
	2 Odilat 3 1320			
Name and	d mailing address of the ISA European Patent Office, P.B. 5818 Patentiaan 2	Authorized officer		
	NL - 2280 HV Ripswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,	0.000		
	Far (+ 31-70) 340-3016	Griffith, G		

Form PCT/ISA/218 (second sheet) (July 1992)

2

INTERNATIONAL SEARCH REPORT

Int. onal Application No PCT/GB 95/02085

		PC1/GB 93/02003
(Continu	non) DOCUMENTS CONSIDERED TO BE RELEVANT	Relevant to claim No.
ategory *	Citation of document, with indication, where appropriate, of the relevant passages	
	CHEMICAL ABSTRACTS, vol. 110, no. 15, 10 April 1989 Columbus, Ohio, US; abstract no. 131899, D. M. BYERS. 'Elongation of exogenous fatty acids by the bioluminescent bacterium Vibrio harveyi.' page 402; column 1; see abstract & J. BACTERIOL., vol. 171, no. 1, pages 59-64,	1-7
A	EP,A,O 309 429 (LIFE SCIENCE INTERNATIONAL AB.) 29 March 1989 see the whole document	1-7
A	FR,A,2 699 930 (P. KABORE) 1 July 1994	
A	US,A,3 666 631 (E. RICH ET AL.) 30 May 1972	
A	US,A,3 423 290 (R. C. SEAMANS, JR. ET AL.) 21 January 1969	

INTERNATIONAL SEARCH REPORT

iformation on patent family members

Ink Shall Application No PCT/GB 95/02085

Patent document cited in search report	Publication date	Patent family member(s)		Publication date	
EP-A-309429		AU-B- SE-A- WO-A-	2484988 8703675 8902929	18-04-89 24-03-89 06-04-89	
FR-A-2699930	01-07-94	NONE			
.US-A-3666631	30-05-72	NONE		****	
US-A-3423290	21-01-69	NONE			

Form PCT/ISA/218 (perent family annex) (July 1992)